

Naval Dental Research Institute

DRI-PR 90-06
November 1990




AD-A231 709

DETECTION OF MICROBIAL TRYPSIN-LIKE ENZYMES BY USE OF AN AGAR GEL

E. D. PEDERSON

B. L. LAMBERTS



DTIC
ELECTE
FEB 07 1991
S B D

Naval Medical Research and Development Command
Bethesda, Maryland

91 2 04 206

NAVAL DENTAL RESEARCH INSTITUTE
BUILDING 1-H
GREAT LAKES, ILLINOIS 60088-5259

DETECTION OF MICROBIAL TRYPSIN-LIKE ENZYMES
BY USE OF AN AGAR GEL

E. D. PEDERSON


B. L. LAMBERTS

Research Progress Report NDRI-PR-90-06
Work Units MR012002-0045 and BS10.DA-440-0
Naval Medical Research and Development Command
National Naval Medical Center
Bethesda, Maryland 20889-5044

The opinions expressed herein are those of the authors and cannot be construed as reflecting the views of the Navy Department or the Naval Service at large. The use of commercially available products does not imply endorsement of these products or preference to other similar products on the market.

This document has been approved for public release; its distribution is unlimited.

Approved and released by



J. C. CECIL III
Captain, Dental Corps
United States Navy
Commanding Officer

Detection of microbial trypsin-like enzymes by use of an agar gel

E. D. Pederson and B. L. Lamberts

Naval Dental Research Institute, Building 1-H, Great Lakes, Illinois 60088-5259, U.S.A.

Abstract

Certain oral micro-organisms associated with periodontal diseases elaborate trypsin-like enzymes. This study describes a simple assay for these enzymes, based on the incorporation of the synthetic substrate N- α -benzoyl-DL-arginine-p-nitroanilide (BAPNA) into ion-agar gel. The system can provide quantitative or qualitative data within 24 h and is suitable for use in the dental clinic.

Introduction

The progression of destructive periodontal diseases may be significantly affected by oral micro-organisms such as black-pigmented *Bacteroides* species and treponemes. *Bacteroides* species have been recovered in high proportions from patients with advanced periodontitis (Slots, 1979; Tanner *et al.*, 1979). Positive correlations have been established for pocket depth, connective tissue attachment loss, and periodontal indexes in patients with a prevalence of subgingival treponemes (Listgarten and Levin, 1981; Armitage *et al.*, 1982). *Treponema denticola* is a specific spirochete which has been shown to be present in elevated numbers in severe periodontitis (Simonson *et al.*, 1988).

A common characteristic of these micro-organisms, which may enhance their virulence, is that they elaborate trypsin-like enzymes (Slots, 1981; Laughon *et al.*, 1982a). This has also been observed for certain species of *Capnocytophaga*, which are often associated with early periodontitis and have been reported to cause neutrophil and fibroblast dysfunctions *in vitro* (Shurin *et al.*, 1979; Stevens *et al.*, 1980). The trypsin-like enzymes were determined by use of the substrate N-benzoyl-DL-arginine-2-naphthylamide (BANA), a component of the API ZYM system (Analytab Products Inc., Plainview, New York, U.S.A.), which provides a semiquantitative micromethod for measuring nineteen enzymatic reactions. This substrate has been employed qualitatively to identify oral *Bacteroides* spp., *Capnocytophaga* spp. and spirochetes (Laughon *et al.*, 1982b). Recently, Tippet *et al.* (1987) described a method for quantitation of *B. gingivalis* (*Porphyromonas gingivalis*) by counting colonies imprinted on filter paper where their trypsin-like enzymatic activities were revealed by use of a BANA-containing chromogenic system.

Another quantitative laboratory procedure for measuring trypsin-like activity has been developed by Haverback *et al.* (1960) by use of the synthetic substrate, N- α -benzoyl-DL-arginine-p-nitroanilide (BAPNA). One advantage of this substrate is that the development of colour from enzymatic action can be monitored during the incubation of a sample. This contrasts with the use of BANA, with which colour development requires the addition of reagents that stop enzymatic action. Preliminary tests of the Haverback BAPNA procedure

at this laboratory, conducted on twelve strains of *B. gingivalis*, confirmed the sensitivity and utility of the method. We demonstrated that elaboration of trypsin-like activity *vs* time of growth peaked at 48–72 h (Lamberts *et al.*, 1985). However, this laboratory procedure was not convenient for clinical use because of the relatively short shelf life of the substrate in solution, the need for additional reagents and the attention required by procedural details.

The present report describes a simple quantitative BAPNA-in-agar assay which can be employed for measuring trypsin-like activities of oral micro-organisms or applied clinically for tests of human plaque samples. Information from the clinical tests could help to identify patients undergoing changes in disease status or could be useful in monitoring the success of therapeutic measures.

Materials and methods

Preparation of BAPNA-in-agar system

A mixture of 0.10–0.15% BAPNA and 1.0% ionagar was dissolved with continuous stirring at 100°C in 0.1 M Tris HCL buffer, pH 7.7. The temperature was adjusted to 60–65°C and aliquots of the mixture were transferred to vials or microplates and permitted to cool to form gels, which were covered with the buffer to prevent dehydration. The gel containers were sealed with lids or plastic wrap and maintained under refrigeration prior to use.

Micro-organisms and culture conditions

Table 1 lists alphabetically the organisms which were tested for their abilities to elaborate trypsin-like enzyme activity. The bacteria were grown to late log or early stationary phases prior to the tests. The following organisms were grown anaerobically for 72 h in Wilkins-Chalgren broth at 37°C: *Bacteroides asaccharolyticus*, *Bacteroides denticola*, *Bacteroides gingivalis*, *Bacteroides intermedius*, *Bacteroides loeschei*, *Bacteroides macacae*, *Capnocytophaga ochracea* and *Capnocytophaga sputigena*.

Application of samples

Pancreatic trypsin III (Sigma) was dissolved in 0.1 M Tris HCL buffer, pH 7.7, (1 mg/ml dry weight as received). This stock solution was employed to prepare 1.25–1,000 ng/100 µl trypsin standard solutions, which were added to microplate wells containing the BAPNA-in-agar. Plates were read at 405 nm immediately after application (blanks) and following suitable incubation periods.

Micro-organisms were applied to the gels either in culture media or as washed cells, suspended in the Tris HCL buffer or in 0.9% saline. Subgingival plaque samples were obtained from patients by curettes and applied to the gels, either directly or after preliminary suspension in saline.

Results

Linear relationships were found between colour development and the amount of pancreatic trypsin, following 37°C incubations of trypsin in the BAPNA-in-agar system (Figure 1). Extending incubations from 2 to 4 or 6 h increased the sensitivity of measurements while retaining the linearity of the plots of activity




Figure 1 Trypsin standard plots following incubation at 37°C. The incubation times were: 2 h, correlation coefficient 0.997; 4 h, correlation coefficient 0.986; and 6 h, correlation coefficient 0.998.

amount of trypsin. Table 1 shows data on trypsin-like activities for sixty-two isolates of oral bacterial species along with information on their Gram-staining properties. It is evident that most of the activity was found with the *B. gingivalis* and *T. denticola* strains. No activity was detectable with the Gram-positive microorganisms.

Tests of twenty-nine subgingival plaque samples revealed a rough correlation of spirochete level and trypsin-like activity. Twenty samples with high to moderately low spirochete levels showed trypsin-like activity, while nine samples with few if any spirochetes showed no activity.

Discussion

Exploratory tests with the subgingival plaque samples indicated that the BAPNA-in-agar system can serve as a rapid, simple method for detecting microbial trypsin-like activity. The system has several potential advantages for clinical use. (a) Plaque samples can be applied directly to a gel by curette, although precautions must be taken to avoid transfer of gel and buffer to the patient's mouth. This may involve the use of a clean curette for each site sampled. (b) Qualitative or semi-quantitative determinations of activity can be made visually. Representative standards may be prepared in vials by incubating various amounts of trypsin with 0.5–2.0 ml volumes of BAPNA-in-agar, then adding 100 μ l of 0.1 N HCl

167 Assay for trypsin-like enzymes

Availability Codes

1st Avail and/or
Special

A-1 20

Table 1 Detectability of trypsin-like activities in oral micro-organisms by BAPNA-in-agar assay

Micro-organism	Gram stain*	Isolate	BAPNA activity
<i>Actinobacillus actinomycetemcomitans</i>	N	ATCC 29523 ⁵	—
		ATCC 29524 ⁵	—
<i>Actinomyces naeslundii</i>	P	ATCC 12104 ⁵	—
<i>Actinomyces odontolyticus</i>	P	ATCC 17982 ⁵	—
<i>Actinomyces viscosus</i>	P	M 100 ⁷	—
<i>Bacteroides asaccharolyticus</i>	N	ATCC 25260 ⁵	—
<i>Bacteroides denticola</i>	N	ATCC 33185 ⁵	+
<i>Bacteroides gingivalis</i> (<i>Porphyromonas gingivalis</i>)	N	D5529 A ¹	+
		D13B 11 ⁴	+
		D40C 4 ¹	+
		D67D 9 ¹	+
		D43A 5 ¹	+
		D82F 5 ¹	+
		376 ¹	+
		2A ¹	+
		ATCC 33277 ⁵	+
		38 ¹	+
		JKG 5 ²	+
		JKG 1 ²	+
		D83T 3 ¹	+
		3B ¹	+
		D868B 6 ¹	+
		D84D 2 ¹	+
		JKG 9 ²	+
<i>Bacteroides intermedius</i>	N	ATCC 25611 ⁵	—
		13043 ¹	—
<i>Bacteroides loescheii</i>	N	ATCC 15930 ⁵	—
<i>Bacteroides macacae</i>	N	ATCC 33141 ⁵	+
<i>Capnocytophaga ochracea</i>	N	ATCC 27872 ⁵	—
<i>Capnocytophaga sputigena</i>	N	ATCC 33612 ⁵	+
<i>Fusobacterium nucleatum</i>	N	ATCC 25586 ⁵	—
<i>Fusobacterium</i> spp.	N	B ¹	—
		5 ¹	—

Table 1 (continued)

Micro-organism	Gram stain*	Isolate	^P BANA activity
<i>Haemophilus aphrophilus</i>	N	ATCC 13252 ⁵	—
<i>Lactobacillus casei</i>	P	1 ³	—
		3 ³	—
<i>Leptotrichia buccalis</i>	N	ATCC 14201 ⁵	—
		ATCC 19616 ⁵	—
<i>Pseudomonas aeruginosa</i>	N	1 ³	—
<i>Proteus vulgaris</i>	N	1 ³	—
<i>Staphylococcus aureus</i>	P	ATCC 12598 ⁵	—
<i>Streptococcus faecalis</i>	P	539 ⁶	—
<i>Streptococcus mutans</i>	P	NCTC 10449 ⁸	—
		6715 ⁹	—
		OMZ 176	—
		C-211	—
<i>Streptococcus sanguis</i>	P	ATCC 10557 ⁵	—
		ATCC 10558 ⁵	—
		410	—
		Challis	—
<i>Treponema denticola</i>	N	ATCC 33520 ⁵	+
		D39DP 1 ⁴	+
		N 39 ⁴	+
		IPP ⁴	+
		Ichelson 2 ⁴	+
		TRRD ⁴	+
		TD 2 ⁴	+
		Ambigua ⁴	+
		D65BR 1 ⁴	+
<i>Treponema vincentii</i>	N	ATCC 33580 ⁵	—

* N, Gram-negative; P, Gram-positive.

¹Forsyth Dental Center; ²University of Michigan; ³Naval Dental Research Institute; ⁴Virginia Polytechnic Institute and State University; ⁵American Type Culture Collection; ⁶University of Notre Dame; ⁷University of West Virginia; ⁸National Culture Type Collection; and ⁹National Institute of Dental Research.

to stop the reaction and stabilize colour. (c) Results with plaque or microbial samples can be obtained within 24–48 h. This procedure may also be useful to elucidate the relationship of plaque microbial composition to periodontal disease status.

Data shown in Table 1 agree with previous reports, in which the substrate BAPNA was used to screen for trypsin-like enzymes from *B. gingivalis* and *T. denticola* (Slots, 1981; Laughon *et al.*, 1982a). Further study of pure cultures may provide comparative information on the stabilities and cell-binding properties of trypsin-like enzymes from various micro-organisms.

The BAPNA-in-agar system has demonstrated a shelf life of over 1 year if maintained frozen or shielded from light at 1–4°C, and of over 3 months at room temperature when shielded from light.

Acknowledgements

The authors wish to thank Dr Lloyd G. Simonson for providing the micro-organisms used in this study, and Ms Myra Portis for manuscript preparation. The opinions expressed herein are those of the authors and cannot be construed as reflecting the views of the Navy Department or the Naval Service at large. The use of commercially available products does not imply endorsement of the products or preference to other similar products on the market. This work was supported by Naval Medical Research and Development Command Project numbers MR012002-0045 and BS10.DA-440-0.

References

- ARMITAGE G. C., Dickinson W. R., Jenderseck R. S., Levine S. M. and Chambers D. W. 1982. Relationship between the percentage of subgingival spirochetes and the severity of periodontal disease. *J. Periodontol.* 53 550–6.
- HAVERBACK B. J., Dyce B., Bundy H. and Edmondson H. A. 1960. Trypsin, trypsinogen and trypsin inhibitor in human pancreatic juice. *Am. J. Med.* 29 424–33.
- LAMBERTS B. L., Shklar I. L., Pederson E. D. and Bial J. J. 1985. Trypsin-like activities of *Bacteroides gingivalis* during 7-day culture periods. *J. Dent. Res.* 64(A) 256.
- LAUGHON B. E., Syed A. A. and Loesche W. J. 1982a. API ZYM system for identification of *Bacteroides* spp., *Campylobacter* spp., and spirochetes of oral origin. *J. clin. Microbiol.* 15 97–102.
- LAUGHON B. E., Syed S. A. and Loesche W. J. 1982b. Rapid identification of *Bacteroides gingivalis*. *J. clin. Microbiol.* 15 345–6.
- LISTERGARTEN M. A. and Levin S. 1981. Positive correlation between the proportions of subgingival spirochetes and motile bacteria and susceptibility of human subjects to periodontal deterioration. *J. clin. Periodont.* 8 122–38.
- SHURIN S. B., Socransky S. S., Sweeney E. and Stossel T. P. 1979. A neutrophil disorder induced by *Campylobacter*, a dental microorganism. *N. Engl. J. Med.* 301 849–54.
- SIMONSON L. G., Goodman C. H., Bial J. J. and Morton H. E. 1988. Quantitative relationship of *Treponema denticola* to severity of periodontal disease. *Infect. Immun.* 56 728–8.
- SLOTS J. 1979. Subgingival microflora and periodontal disease. *J. clin. Periodont.* 6 351–82.
- SLOTS J. 1981. Enzymatic characterization of some oral and nonoral Gram-negative bacteria with the API ZYM system. *J. clin. Periodont.* 14 288–94.
- STEVENS R. H., Sela M. N., Shapira J. and Hammond B. F. 1980. Detection of a fibroblast proliferation inhibitory factor from *Campylobacter sputigena*. *Infect. Immun.* 27 271–5.

TANNER A. C. R., Haffer C., Bratthall G. T., Visconti R. A. and Socransky S. S. 1979. A study of the bacteria associated with advancing periodontitis in man. *J. clin. Periodont.* 6 278–307.

TIPPET B., Frisken K. W. and Tagg J. R. 1987. Quantification of the bacterium *Bacteroides gingivalis* in human dental plaque by detection of the trypsin-like protease activity of colonies imprinted on membrane filters. *Archs Oral Biol.* 32 151–2.

Accepted 5 March 1990


UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) DETECTION OF MICROBIAL TRYPSIN-LIKE ENZYMES BY USE OF AN AGAR GEL		5. TYPE OF REPORT & PERIOD COVERED
7. AUTHOR(s) E. D. PEDERSON and B. L. LAMBERTS		6. PERFORMING ORG. REPORT NUMBER NDRI-PR 90-06
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Dental Research Institute Building 1-H Great Lakes, IL 60088-5259		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Medical Research and Development Command, National Naval Medical Center, Bethesda, MD 20889-5044		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS MR012002-0045 and BS10.DA-440-0
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Chief, Bureau of Medicine and Surgery Navy Department Washington, D.C. 20372-5120		12. REPORT DATE November 1990
		13. NUMBER OF PAGES 7
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release; distribution unlimited.		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) This document has been approved for public release; distribution unlimited.		
18. SUPPLEMENTARY NOTES Microbios 63:165-171, 1990		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Bacterial enzyme BAPNA Synthetic substrate Trypsin-like enzymes		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Certain oral micro-organisms associated with periodontal diseases elaborate trypsin-like enzymes. This study describes a simple assay for these enzymes, based on the incorporation of the synthetic substrate N- α -benzoyl-DL-arginine-p-nitroanilide (BAPNA) into ion-agar gel. The system can provide quantitative or qualitative data within 24 h and is suitable for use in the dental clinic.		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)



S N 0102- LF- 014- 6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)